

# Mechanistic investigation of a novel medium-sized ring rearrangement

Al-Ani, Aneesa; Jones, Alan M.

DOI:

[10.1055/s-0036-1589141](https://doi.org/10.1055/s-0036-1589141)

License:

Other (please specify with Rights Statement)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Al-Ani, A & Jones, AM 2018, 'Mechanistic investigation of a novel medium-sized ring rearrangement', *Synlett*, vol. 29, no. 05, pp. 589-592. <https://doi.org/10.1055/s-0036-1589141>

[Link to publication on Research at Birmingham portal](#)

## **Publisher Rights Statement:**

Checked for eligibility: 13/04/2018

© 2018 Georg Thieme Verlag KG

<https://www.thieme-connect.de/DOI/DOI?10.1055/s-0036-1589141>

## **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.


# Mechanistic Investigation of a Novel Medium-sized Ring Rearrangement

Aneesa J. Al-Ani<sup>a</sup> and  
Alan M. Jones<sup>b\*</sup>

<sup>a</sup> Division of Chemistry and Environmental Science, Manchester Metropolitan University, M1 5GD, UK

<sup>b</sup> School of Pharmacy, University of Birmingham, Edgbaston, B15 2TT, UK


\* Corresponding author: Dr Alan M. Jones, email  
[a.m.jones.2@bham.ac.uk](mailto:a.m.jones.2@bham.ac.uk) phone +44(0)1214-147288, web:  
[www.jonesgroupresearch.wordpress.com](http://www.jonesgroupresearch.wordpress.com)



Received:  
Accepted:  
Published online:  
DOI:

**Abstract** The dominant mechanism operating in a recently disclosed ring-contraction is revealed for the first time. Providing new insight and supporting evidence into a classic ring contraction strategy within a rigidified larger ring system.

**Key words** Favorskii, mechanism, medium-sized ring, <sup>18</sup>O labelling, Rearrangement, Ring contraction, Semi-benzylic.




**Scheme 1** Observed ring contraction of 1 to 2 and isolated intermediate 3.

One of the elegant solutions in the organic chemists' toolbox, when constructing a densely functionalised smaller ring system, for purposes as varied as natural product synthesis<sup>1</sup> to medicinal chemistry,<sup>2</sup> is to contract an easier to access larger ring system.<sup>3</sup> Ring contractions can proceed *via* one of three dominant mechanisms: cationic, carbenoid or anionic in nature.<sup>4</sup> The importance of this class can easily be seen with named reactions such as the Pinacol, semi-Pinacol, Benzilic acid, semi-Benzylic and Favorskii rearrangements.<sup>5</sup> Recently, we identified within a class of 10-membered heterocyclic ring systems (7,8,9-trihydrobenzo[*d*]quinazolo[1,2,3-*a,b*,1,3]diazecane-1,10,11,17-tetrones (1))<sup>6</sup> a one-pot, multi-step rearrangement under mild conditions to pyrrolo[2,3-*b*]quinolines (such as 2, Scheme 1) exemplified with alkoxides and amines in up to 77% yield.<sup>7</sup> Fortuitously, a competent isolated intermediate (3) was found in certain examples, that when re-submitted to the reaction conditions for a further length of time afforded the same product as 2. We previously hypothesised that it was likely either a benzylic or Favorskii-type rearrangement occurred from reaction intermediate (3) that led to the ring contraction product (2).

In Scheme 1 it can be seen that isolated intermediate 3 can exist in equilibria with its diketone tautomeric structure (A). Pathway A, depicts a quasi-Favorskii rearrangement proceeding *via* cyclisation of the N(10) enamine of A onto the C(5) carbonyl group will form a strained cyclopropanone (intermediate B). The cyclopropanone (B) will be labile to attack by a nucleophile and in doing so the ring strain will be relieved. Elimination of hydroxide from B will in turn generate an aromatic quinoline ring. An equally likely route to 2 from 1 is Pathway B, a semi-benzylic rearrangement of A due to the non-enolizable diketone functionality present and in particular, the highly electrophilic C5 carbonyl group (δ 178.3 ppm, <sup>13</sup>C NMR). Therefore, attack of a nucleophile at the C5 position of A could induce a ring contraction via intermediate C to 2.<sup>7</sup> Herein we show for the first time, using isotopic labelling studies, the major mechanism in operation.

To conclusively prove whether either of these rearrangements (pathway A or B, *quasi*-Favorskii or semi-benzylic reactions, respectively) occurred we decided upon a microscale <sup>18</sup>Oxygen-labelling experiment of the carbonyl group's installed via oxidative fragmentation in the preceding step (Scheme 2).



**Scheme 2** Potential ring-contraction mechanisms in operation. Key: sB = semi-Benzylic; qF = quasi-Favorskii.


This step would determine if the  $^{18}\text{O}$ -enriched C(4) carbonyl oxygen was retained or lost during the ring contraction step. In Scheme 2 it can be seen that labelling the C(4) carbonyl of intermediate **3** (and its tautomer **A**) would yield binary results. The alternative incorporation of a  $^{13}\text{C}$  label from a reaction precursor would be retained in both rearrangement route products (not depicted).<sup>8</sup>

Operative mechanisms (Scheme 2) include two routes to a semi-benzylic mediated ring contraction and two routes to a quasi-Favorskii mediated ring contraction, respectively. Addressing the four potential routes in turn; the *quasi-Favorskii* route A1 which would proceed with retention of aromaticity and retention of label and route A2 via an anti-aromatic intermediate and loss of label. In turn, *semi-Benzylic* route B1 mechanism proceeds through nucleophilic attack at the more electrophilic C5 carbonyl of **3** (178.3 ppm,  $^{13}\text{C}$  NMR). Whilst *semi-Benzylic* route B2 mechanism proceeds through nucleophilic attack at the less electrophilic C4 carbonyl of **3** (153.1 ppm,  $^{13}\text{C}$  NMR). Comparison of route B1 with route B2 would lead to different outcomes, loss of label (route 1) and retention of label (route 2), respectively. Therefore, if the  $^{18}\text{O}$  label was retained a *quasi-Favorskii* mechanism (route A1) would be the most likely mechanism in operation (due to ground state carbonyl electrophilicity measurements) and loss of label would be most likely due to a *semi-Benzylic* rearrangement (Route B1, mitigating the need to undergo a strained anti-aromatic intermediate). The further possibility on an internal  $6\pi$  electrocyclisation (*c.f.* Karimi)<sup>9</sup> of intermediate species **3** was ruled out due to the internal strain within the multi-ring fused system.

Due to the scarcity of  $^{18}\text{O}$ -labelling reagents, a microscale experiment was chosen to prepare  $^{18}\text{O}$ -*m*-CPBA from commercially available (90% labelled)  $^{18}\text{O}$ -hydrogen peroxide (Scheme 3).<sup>10</sup> Briefly, fusion of methyl anthranilate with  $\delta$ -valerolactam in a sealed reactor vessel afforded **4** in 41% yield.<sup>11</sup> Compound **4** was exposed to a single equivalent of freshly prepared  $^{18}\text{O}$ -*m*-CPBA under high-dilution conditions, until consumption of **4** was indicated by TLC. The intermediate species was treated with a further equivalent of unlabelled *m*-CPBA to ensure all of the  $^{18}\text{O}$ -labelled epoxide was intercepted to undergo Grob-type fragmentation to **1** in a 40% isolated yield. The location of the  $^{18}\text{O}$  carbonyl group can be inferred from the well-known nitrogen assisted ring opening of epoxides from previously reported trapping experiments.<sup>12</sup> Furthermore, ground state measurement of the infrared carbonyl absorption


frequency would be expected to undergo a subtle change. Due to the overlapping of the diketone absorption, the discrimination of the two carbonyl groups was not possible. However, a subtle difference in the absorption frequency of the diketone unit (1707  $\text{cm}^{-1}$  in the labelled versus 1709  $\text{cm}^{-1}$  in the unlabelled infrared spectra) and no change in the urea/amide carbonyl stretching frequency gave further credence to the location of the  $^{18}\text{O}$  carbonyl location being on the newly installed carbonyl of the diketone in **1** versus the urea carbonyl, in accordance with the proposed reaction mechanism.

$^1\text{H}$  and  $^{13}\text{C}$  NMR spectroscopic analyses were in accordance with the previously reported structure of unlabelled **1**,<sup>6</sup> furthermore electrospray mass spectroscopic analyses indicated a mono-incorporation of an  $^{18}\text{O}$ -label with slight loss of labelling.<sup>13</sup>



**Scheme 3** Preparation of  $^{18}\text{O}$ -labelled reaction precursor **1** and mechanistic rationale.

$^{18}\text{O}$ -labelled **1** was treated<sup>14</sup> under the reported conditions of methanolic sodium methoxide to afford the ring contraction product **2** in a 75% isolated yield, as confirmed by NMR spectral analyses (Scheme 4).



**Scheme 4** Treatment of **1** with NaOMe could afford labelled-**2** or unlabelled-**2**

Analysis of the mass spectrum of **2** revealed no trace of the  $^{18}\text{O}$ -label presence (other than at background levels) indicating **2** was unlabelled. Therefore, based on the possible reaction pathways outlined in Schemes 1 and 2, the predominant pathway in operation was *semi-benzylic*, in agreement with avoiding a strained anti-aromatic reaction intermediate (Scheme 2) and being driven by nucleophilic attack at the more electrophilic carbonyl. Further supporting evidence for this identified reaction pathway can be observed by the migratory aptitude trend of a phenyl group over a carbonyl group migration in related systems.<sup>15</sup>

In this letter we have identified the predominant mechanism of ring contraction within a 10-membered ring system (via a 7,5-ring system) to a 6,5-ring system is *semi-benzylic* in nature through  $^{18}\text{O}$ -isotopic labelling studies. Providing new insight into ring contraction mechanisms within complex substructures.

## Funding Information

HEFCE

## Acknowledgment

Dr Maira Guzman (Manchester Metropolitan University) is thanked for mass spectroscopic analyses.

## Supporting Information

YES (this text will be updated with links prior to publication)

## Primary Data

NO (this text will be deleted prior to publication)

## References and Notes

- (1) (a) Silva, L. F. **2013** Ring Contraction Reactions in the Total Synthesis of Biologically Active Natural Products. *Stereoselective Synthesis of Drugs and Natural Products*, Wiley, 1-20; (b) Sun, C.; Lee, H.; Lee, D. *Org. Lett.* **2015**, *15*, 5348-5351; (c) Craig, D.; Funai, K.; Gore, S. J.; Kang, A.; Mayweg, A. V. W. *Org. Biomol. Chem.* **2011**, *9*, 8000-8002.
- (2) Bull, J. A.; Croft, R. A.; Davis, O. A.; Doran, R.; Morgan, K. F. *Chem. Rev.* **2016**, *116*, 12150-12233.
- (3) Hong, A. Y.; Krout, M. R.; Jensen, T.; Bennett, N. B.; Harned, A. M.; Stoltz, B. M. *Angew. Chem. Int. Ed.* **2011**, *50*, 2756-2760.
- (4) Silva, L. F. *Tetrahedron* **2002**, *58*, 9137-9161.
- (5) (a) Grainger, R. S.; Betou, M.; Male, L.; Pitak, M. B.; Coles, S. J. *Org. Lett.* **2012**, *14*, 2234-2237; (b) Song, Z.-L.; Fan, C.-A.; Tu, Y.-Q. *Chem. Rev.* **2011**, *111*, 7523-7556; (c) Hamon, D. P. G.; Tuck, K. L. *Chem. Commun.* **1997**, 941-942; (d) Lee, E.; Yoon, C. H. *J. Chem. Soc., Chem. Commun.* **1994**, 479-481.
- (6) Jones, A. M.; Liu, G.; Lorion, M. M.; Patterson, S.; Lebl, T.; Slawin, A. M. Z.; Westwood, N. J. *Chem. Eur. J.* **2011**, *17*, 5714-5718.
- (7) Jones, A. M.; Patterson, S.; Lorion, M. M.; Slawin, A. M. Z.; Westwood, N. J. *Org. Biomol. Chem.* **2016**, *14*, 8998-9011.
- (8) Denmark, S. E.; Venkatraman, S. J. *Org. Chem.* **2006**, *71*, 1668-1676.
- (9) Karimi, S.; Ma, S.; Ramig, K.; Greer, E. M.; Szalda, D. J.; Subramaniam, G. *Tetrahedron Lett.* **2015**, *56*, 6886-6889.
- (10) (a) McDonald, R. N.; Steppel, R. N.; Dorsey, J. E. *Org. Synth.* **1988**, *6*, 276; (b) Jankowski, S.; Kaminski, R. *J. Label Compd. Radiopharm.* **1995**, *36*, 373-376; (c) Steyn, P. S.; Vleggaar, R. J. *Chem. Soc., Chem. Commun.* **1985**, 1531-1532.
- (11) (a) Patterson, S.; Lorenz, C.; Slawin, A. M. Z.; Westwood, N. J. *QSAR & Comb. Sci.* **2004**, *23*, 883-890; (b) Moehrle, H.; Seidel, C. *Chem. Ber.* **1973**, *106*, 1595-1600.
- (12) (a) Davies, S. G.; Key, M. S.; Rodriguez-Solla, H.; Sanganee, H. J.; Savory, E. D.; Smith, A. D. *Synlett* **2003**, 1659-1662; (b) Aciro, C.; Davies, S. G.; Garner, A. C.; Ishii, Y.; Key, M.-S.; Ling, K. B.; Prasad, R. S.; Roberts, P. M.; Rodriguez-Solla, H.; O'Leary-Steel, C.; Russell, A. J.; Sanganee, H. J.; Savory, E. D.; Smith, A. D.; Thomson, J. E., *Tetrahedron* **2008**, *64*, 9320-9344.
- (13) See electronic supporting information for further details and Gardner Swain, C.; Tsuchihashi, G.-I.; Taylor, L. J. *Anal. Chem.* **1963**, *35*, 1415-1417.
- (14) Typical procedure: To a solution of **1** (10 mg, 0.03 mmol) in anhydrous methanol (1.0 mL) was added sodium methoxide (3 mg, 0.06 mmol). The reaction mixture was stirred at room temperature for 16 h then evaporated *in vacuo*, re-dissolved in dichloromethane (4.0 mL) and washed with NaHCO<sub>3</sub>(aq.) (2 × 2.0 mL). The organic layer was dried (MgSO<sub>4</sub>), filtered and concentrated *in vacuo* to give a solid which was purified by flash column chromatography (SiO<sub>2</sub>; ethyl acetate:hexane (3:7)) furnished to afford the title compound as a yellow powder (8 mg, 75%). Mp 148-149 °C (lit.<sup>S3</sup> 149-150 °C); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>): δ 8.18 (dd, <sup>3</sup>J = 8.5 Hz, <sup>4</sup>J = 1.0 Hz, 1H), 7.88 (dd, <sup>3</sup>J = 8.0 Hz, <sup>4</sup>J = 1.5 Hz, 1H), 7.63-7.52 (m, 2H), 7.49-7.42 (m, 1H), 7.36-7.31 (m, 1H), 7.30-7.23 (m, 2H), 4.20 (t, <sup>3</sup>J = 8.0 Hz, 2H), 4.04 (s, 3H), 3.55-3.47 (m, 5H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>): δ 168.0, 166.8, 158.3, 147.6, 139.2, 132.3, 131.2, 130.7, 128.9, 128.6, 127.0, 126.9, 124.9, 124.8, 123.4, 122.9, 120.9, 52.3, 51.9, 50.3, 26.9; IR (KBr): ν<sub>max</sub> = 2945 (m), 1725 (s) (C=O), 1717 (s) (C=O), 1617 (m), 1597 (m), 1483 (m), 1228 (w), 754 (w) cm<sup>-1</sup>; LRMS (ESI<sup>+</sup>): m/z (%) 363 [M+H]<sup>+</sup>; HRMS (ESI<sup>+</sup>): m/z calcd for C<sub>21</sub>H<sub>19</sub>N<sub>2</sub>O<sub>4</sub> [M+H]<sup>+</sup>: 363.1345; found 363.1341.
- (15) Nakamura, K.; Osamura, Y. *J. Am. Chem. Soc.* **1993**, *115*, 9112-9120.

## Checklist (have these on hand for manuscript submission in ScholarOne):

- cover letter, including a statement of the work's significance
- full mailing address, telephone and fax numbers, and e-mail address of the corresponding author
- email address for each author
- original Word file
- original graphics files zipped into one zip file
- eye-catching graphical abstract as an individual file
- 5-8 key words
- separate Supporting Information file
- separate zipped Primary Data files including cover sheet (optional)

## Useful links:

- [SYNLETT homepage](#)
- [SYNLETT information and tools for authors](#)
- [Graphical abstract samples](#) (PDF file download)
- [What is "Primary Data"?](#)
- [ScholarOne](#) (manuscript submission)